



## WAVE 4

National Income Dynamics  
Study (NIDS) – Coronavirus  
Rapid Mobile Survey (CRAM)

# COVID19 Vaccine Intentions in South Africa: Health communication strategy to address vaccine hesitancy

Uma Kollamparambil - University of the Witwatersrand


Adeola Oyenubi - University of the Witwatersrand

Chijioke O. Nwosu - Human Sciences Research Council

12 May 2021



**N.i.D.S.**  
NATIONAL INCOME DYNAMICS STUDY



CORONAVIRUS RAPID MOBILE SURVEY 2020

# COVID19 Vaccine Intentions in South Africa: Health communication strategy to address vaccine hesitancy

**Uma Kollamparambil**, School of Economics & Finance, University of the Witwatersrand<sup>1</sup>.

**Adeola Oyenubi**, School of Economics & Finance, University of the Witwatersrand<sup>2</sup>.

**Chijioke O. Nwosu**, The Impact Centre, Human Sciences Research Council, South Africa<sup>3</sup>.

12 May 2021

## Abstract

Widespread vaccine acceptance is key to achieving herd immunity through vaccination against COVID19, especially because the available vaccines do not have 100% efficacy. The study finds that vaccine acceptance is lower than that of non-pharmaceutical intervention like wearing facemask in public. Based on the vaccine intentions of the respondents of National Income Dynamics Study-Coronavirus Rapid Mobile (NIDS-CRAM) survey, only 55% are fully accepting of the vaccine, while a further 16% are moderately accepting of vaccines. Together, vaccine acceptance is estimated at 70.8% (with 95% confidence interval of 70%-72%) and vaccine hesitancy against COVID19 is estimated at 29.2% (with 95% confidence interval of 28%-30%) amongst the adult South African population. The reasons cited for vaccine hesitancy are primarily due to concerns regarding the side-effects and efficacy of vaccine.

Using the Health Belief model and the Extended Parallel Process model, the study has identified perceived risk of infection with the mediating role of efficacy as a key predictor of vaccine intention. Apart from vulnerability to infection, those perceiving higher severity of risk (elderly and those with chronic illness) have lower hesitancy compared to their younger and healthier counterparts. Higher awareness of COVID19 related information and higher household income are correlated with lower vaccine hesitancy.

There are other significant differences across socio-economic and demographic variables in vaccine hesitancy. The non-black African population group has significantly high vaccine hesitancy compared to black Africans. Males on average have lower hesitancy compared to females, and unmarried/unpartnered individuals are found to have higher hesitancy.

From a communication perspective, it is imperative to continue risk messaging, hand in hand with clearer information on the efficacy of the vaccines. The risk messaging needs to be targeted at younger age and low-income groups who perceive the lowest risk. Those who are less informed on COVID19 are also found to be those who perceive low-risk, therefore this reinforces the need to have in place a communication plan to educate and create awareness on the pandemic. There is an all-around need to enhance vaccine efficacy related messaging, particularly targeting non-blacks and religious individuals. Concerns regarding the side-effects of vaccines need to be addressed with more transparent scientific information together with the associated risk-benefit analysis.

---

<sup>1</sup> [Uma.kollamparambil@wits.ac.za](mailto:Uma.kollamparambil@wits.ac.za)

<sup>2</sup> [Adeola.oyenubi@wits.ac.za](mailto:Adeola.oyenubi@wits.ac.za)

<sup>3</sup> [Cnwosu@hsrc.ac.za](mailto:Cnwosu@hsrc.ac.za)

# Executive Summary

With the focus of government now shifting to making COVID19 vaccines available and its roll-out strategy, it is equally important to work towards ensuring the public uptake of vaccines once they become widely available. While the President of South Africa has announced that vaccination against COVID19 would be purely voluntary, large scale vaccine uptake is required to ensure its effectiveness. This is because the success of the vaccination programme depends upon achieving a scale that is sufficient to result in community immunity, especially given the fact that the COVID19 vaccines do not have 100% efficacy. Further, continued infection amongst the unvaccinated can lead to heightened risk of further virus mutation, exposing even those vaccinated (against the earlier strains of the virus) to new virus strains. Therefore, there are social benefits in minimising vaccine hesitancy.

The objective of this study is to assess COVID19 vaccine intentions in South Africa, identify the socio-economic patterns in vaccine hesitancy and to develop a targeted health communication strategy based on the Health Belief and Extended Parallel Process models in order to improve vaccine acceptance.

The main findings of the study are as follows:

- Based on the responses to the question on vaccine intention, 55% of the population are seen to have strong acceptance of the vaccine. A further 16% are found to accept the vaccine, but with lower certainty. Together, vaccine acceptance is estimated to be 70.8% (with 95% confidence interval of 70%-72%) of adult South African population.
- Commensurately, vaccine hesitancy among adult South African population is estimated as being 29.2% (with 95% confidence interval of 28%-30%). This is a significant proportion of population and raises concerns regarding the success of achieving herd immunity through vaccination.
- Comparing the findings of this study with the meta-analysis by Sallam (2021), vaccine acceptance in South Africa is lower than countries like Ecuador, Malaysia, Indonesia and China; but higher than countries like Kuwait, Jordan, Italy, Russia, Poland, US, and France.
- Vaccine acceptance in South Africa is substantially lower than the non-pharmaceutical behavioural changes undertaken as protective measure against COVID19. The lower acceptance of vaccine as against non-pharmaceutical interventions (NPI) is to be expected considering the perceived dangers of secondary harm of pharmaceutical interventions compared to NPIs. Nonetheless, the results indicate that individuals who take NPIs seriously are also more likely to accept vaccination.
- Among the reasons cited for vaccine hesitancy, the highest proportion cited concerns regarding side-effects of vaccines (30%). This is not surprising considering the lack of public information on vaccine trials as well as emerging media reports on possible serious side-effects of certain vaccines like blood-clotting (CIDRAP 2021).
- Concerns regarding the side-effects of vaccine aside, almost 19.4% of those who indicated unwillingness to accept the COVID19 vaccine pointed to its ineffectiveness as a reason for hesitancy. The mixed messaging coming through media on the efficacy of vaccines as well as their uncertainty regarding the usefulness against the new COVID strains are contributing to the concerns regarding ineffectiveness of vaccine.
- The concerns regarding side-effects and ineffectiveness (jointly accounting for almost 50% of those hesitant) are specific to COVID19 vaccines and therefore needs to be addressed with specific information on COVID19 vaccines. The lack of clarity in communication and unavailability of information on vaccines are clearly contributing to vaccine hesitancy.
- General anti-vaccine stance (that extends beyond COVID specific vaccine inhibition) and lack of trust in the government accounted for nearly 17% of individuals who indicated vaccine hesitancy.

This category of individuals will need to be addressed with information beyond COVID-specific vaccine. The approach taken would have to be broader, but nevertheless COVID specific information will also be useful in generating awareness on vaccine efficacy.

- Severity of risk also is strongly associated with vaccine intention. Those with chronic illness and higher age have lower probability of vaccine hesitancy. This is not surprising given the lower risk perception among the young and the healthy with regards to the severity of COVID19 infection.
- Non-black African population group has significantly high vaccine hesitancy compared to black Africans. Younger adults, females and unmarried/unpartnered individuals are also found to have higher hesitancy. These groups have to be targeted for vaccine related communication.
- The Health behaviour models offer important predictors of vaccine intentions. Risk perception (both vulnerability and severity) is the key driving factor through the multivariate regression analysis. The group that was seen to be most accepting of vaccines are those that perceive high risk. It is therefore important to target risk messaging to the 53% of population that is found to have low risk perception.
- The risk messaging needs to be targeted at younger age population who perceive the lowest risk. Those who are less informed on COVID19 are also found to be those who perceive low-risk, therefore this reinforces the need to put in place a communication plan to educate and create awareness on the pandemic.
- The study findings validate the EPPM construct of the mediating role of efficacy in driving the behavioural response to risk perception. This underscores the need to focus on educating the public about vaccine efficacy to reduce hesitancy.
- It is important to address the reasons that were cited for vaccine hesitation. These include primarily the possible negative side-effects, and ineffectiveness of vaccine. While the risk of blood clots associated in particular with AstraZeneca and Johnson& Johnson vaccines has grabbed headlines, the risk of the vaccine vis-a-vis more accepted pharmaceutical practices like the use of contraceptive pills is considered to be lower (Taylor 2021). Although the survey does not allow pinpointing the nature of side-effect that is causing vaccine hesitancy, it still underlines the need to make available more information from the scientific studies on the possible side-effects and effectiveness that would address possible misinformation circulating through social media and allow individuals to make more informed judgements based on risk-benefit assessments. The lack of information in this regard is contributing substantially to vaccine hesitancy and there is a clear need for better communication around this.
- Increasing vaccine acceptance requires a three-pronged health communication strategy being put in place that targets risk messaging to the young, less educated, less informed, lower income individuals; efficacy messaging to non-black and religious individuals; call to action among older, educated and well-informed individuals.
- South Africa is a religious country and channelling risk as well as efficacy communication through religious leaders is recommended
- The targeted communication aside, there is an over-arching need to provide research-based information on the side-effects and effectiveness of COVID 19 vaccines across the board. Concerns regarding the side-effects of vaccines need to be addressed with more transparent scientific information together with the associated risk-benefit analysis.

# 1. Introduction

While the focus of protective behaviour for the first year of the COVID19 pandemic relied on non-pharmaceutical interventions (NPI), with the successful development of various COVID19 vaccines, the available approaches to protective behaviour in South Africa are expected to widen to include pharmaceutical interventions in the form of vaccines in the near future. Previous studies have shown that acceptance of NPI<sup>4</sup> based protective behaviour is high in South Africa and has improved from 92% in May-June 2020 to 97 % in July-August 2021 as the pandemic progressed (Kollamparambil & Oyenubi 2020, Burger et al 2020). However, the secondary effects of NPI and vaccines can be very different and therefore the behavioural response to both can differ substantially. Hence it is timely to assess the vaccine intentions of the population with a view to put in place measures to improve vaccine acceptance.

The focus of government has now shifted to making vaccines available and its roll-out strategy. It is equally important to work towards ensuring the public uptake of vaccines once they become widely available (Dror et al 2020). While the President of South Africa has announced that vaccination against COVID19 would be purely voluntary, large scale vaccine uptake is required to ensure its effectiveness. This is because the success of the vaccination programme depends upon achieving a scale that is sufficient to result in community immunity. Further, continued infection amongst the unvaccinated can lead to heightened risk of further virus mutation, exposing even those vaccinated (against the earlier strains of the virus) to new virus strains (Rubin 2021). Therefore, there are social benefits in minimising vaccine hesitancy.

Pre-COVID19 studies have acknowledged that behavioural response against vaccination can be specific to a particular vaccine depending on the perceived risk of infection as well as the confidence in the vaccine (Dube et al 2013). Sallam (2021) has highlighted the wide variation between COVID19 vaccine acceptance between countries. It is therefore highly relevant to assess vaccine intentions in South Africa with regard to the COVID vaccine and explore ways to improve vaccine acceptance in the general public health interest.

Various studies have shown that health communication has an important role to play in influencing individual response to health risks (Latimer et al. 2010; Lustria, et al. 2013; Noar et al. 2011). Hence it is imperative to develop a health communication strategy aimed at encouraging individuals to accept both NPI as well as vaccination against COVID19.

While different theories and models have been put forth by health communication theorists designed for health messaging, the Health Belief (Rosenstock 1974) and the Extended Parallel Process (EPP) (Witte 1992; 1994) models are most suited in the pandemic context because of their foundation on fear appeal or health risk (Noar 2004, Dillard, 1994). Meta-analyses of fear appeal messages have found that manipulations of fear appeals have an impact on attitudes, intentions, and behavior changes (Boster & Mongeau, 1984; Witte & Allen, 2000). Therefore, understanding the constructs of risk and efficacy using fear appeal theories such as Health Belief Model (HBM) and Extended Parallel Process model (EPPM) to encourage protective behaviour against COVID19 can be useful.

Health Belief Model (HBM) is one of the earliest health models to use key risk perception variables along with efficacy and barriers to behavioural response to predict health response behaviour. Building on this, the extended parallel process model (EPPM) (Witte, 1992, 1994) posits that efficacy plays a mediating role in the relationship between risk perception and response to the fear appeal message. Based on the interaction between risk perception and perceived efficacy, Witte (1992) identifies two possible outcomes even with high perceived risk (High risk-High efficacy and High risk-Low efficacy) based on the available risk reduction measures. One of the implications of this is recognition that greater fear does not necessarily lead to greater message acceptance, but can perversely cause message rejection, a possibility not acknowledged by HBM. Therefore, identifying the risk-efficacy interaction is relevant in tailoring the communication strategy aimed at improving health response behaviour.

---

4 The most prominent NPI measures include the use of facemask, hand hygiene, social distancing, and staying at home.

The HBM and EPPM frameworks have been used extensively to assess and predict the effect of health messaging on the recommended behavioural responses. Pre-COVID studies have used the EPPM in the context of various health issues such as cardiovascular diseases, hygienic behavior, public health emergency response, teen pregnancy, smoking, vaccination, and HIV/AIDS, among others (Carcioppolo et al., 2013; McKay et al. 2004; Slonim et al., 2005; Witte, 1997; Witte et al. 2002); and also specifically relating to vaccine confidence (Barnett et al 2013; Smith et al 2011).

Studies within the COVID context are fast emerging using the EPPM framework (Lithopoulos 2021; Shirahmadi et al. 2020; Lin and Chen 2020, Yang et al 2020) or the HBM framework (Kollamparambil and Oyenubi 2021). However, these studies relate primarily to non-vaccine behaviour. Exceptions are Chu and Liu (2021) in the US context and Lin et al. (2020) in the context of China which used the EPPM and HBM frameworks respectively to assess COVID19 vaccine behaviour. Edwards et al (2021) has identified the correlates of vaccine hesitancy in Australia, without the use of a specific theoretical framework. The authors have not been able identify a study in the South African context on COVID19 vaccine intention.

The objective of this study is to assess the level of COVID19 vaccine hesitancy in South Africa, identify the socio-economic patterns in vaccine hesitancy and to develop a targeted health communication strategy based on the HBM and EPPM models in order to improve vaccine acceptance.

## 2. Health Belief Model (HBM) and Extended Parallel Process Model (EPPM)

Noar (2004) has highlighted the scarcity of empirical studies that compare the different health behaviour theories in their analysis, with most studies choosing to stick to one selected theory as the framework. This study sees value in comparing the HBM and EPPM. Both these theories are based on fear assessment as the major motivator of behavioural change (Noar 2004) and as such are suited to the pandemic context.

HBM is one of the earliest health behaviour theories developed from 1950s (Rosenstock 1974). The four pillars of HBM model are risk susceptibility/vulnerability, risk severity, efficacy, and barriers to behavioural change. EPPM builds on this foundation using also the tenets of the parallel process model (PPM) (Leventhal 1970) which distinguished between two independent reactions to fear appeals, viz, (a) a cognitive response of risk management process, leading to protective behaviour, and (b) an emotional response of fear management process, that leads to denial, and avoidance. While PPM did not explain the contexts when either of these responses is evoked, EPPM extends the PPM to incorporate the four elements (the perceived severity of a threatening event, the perceived vulnerability, the response efficacy of the recommended preventive behaviour, and the perceived self-efficacy) of Protection Motivation Theory (PMT) (Rogers 1975) to explain this.

How individuals respond to fear appeal is explained by the EPPM using the two components of risk (i.e., severity and vulnerability) and the two components of efficacy (i.e., response efficacy, or the belief in the preventive measure yielding the desirable result and; self-efficacy, or the belief in one's ability to effectively adopt the preventive measure) (Witte, 1992; Witte 1994). According to EPPM, protection motivation is highest when all four cognitions are high leading to the highest level of adaptive change in health attitudes and behaviour. According to the model, the processing of the fear/risk messaging starts with risk appraisal. If the risk appraisal is high, it motivates the individual to start the second appraisal viz., the assessment of efficacy (Witte, 1992).

However, where fear messaging leads to high risk perceptions but the response actions recommended in the message are ineffective, then individuals are incapable of responding appropriately. This leads to the individual responding to the fear and not the danger. Often this situation results in defensive motivation whereby the individual ignores the message or is in denial of the existence of the risk (Witte, 1992). It is therefore clear that while risk perception is important in eliciting a

cognitive response to danger control, it needs to be accompanied by efficacy to ensure appropriate health behaviour response.

According to EPPM, if the individual perceives low risk, then there is no motivation to adopt behavioural change even if the individuals have high efficacy. Lastly, individuals with both low risk and low efficacy are considered to be indifferent as they are unaware of both risk and protective measures. Risk messaging can be considered to be effective when both the risk appraisal as well as efficacy are high as it evokes appropriate danger control process. *Table 1* provides a summary of the interactions between risk and efficacy as outcomes of risk messaging. The appropriate health communication strategy has to be aligned with the response of individuals to risk and efficacy.

**Table 1: EPPM Segments**

	High Efficacy: Belief that infection can be avoided due to the effectiveness of solutions and high feasibility to practice them#	Low Efficacy: Doubt that infection can be avoided due to the perceived ineffectiveness of vaccine and NPI and the low feasibility to adopt the measures#
High Risk: Belief that the risk of infection is high and the danger from infection is high	Responsive: People are educated about risk and respond by taking protective action	Denial: People are aware of the risks but feel helpless to respond effectively.
Low Risk: Belief that the risk of infection is low and the severity of risk from infection is low	Unmotivated: People know what to do but are not really motivated to do much as risk perception is low.	Indifferent: People don't feel at risk and are unaware of prevention strategies.
	Low risk: Due to the low risk perception, people do not enter into efficacy assessment	

**Source:** Adapted from Rimal (2001)

# COVID19 being a communicable disease, the efficacy beliefs depend not just on one's own ability to adopt the measures but also on the ability of others to do so.

While *Table 1* identifies, two categories of individuals with low risk (with high efficacy and low efficacy), Witte (1992) points out that it is irrelevant as the efficacy appraisal does not kick-in for an individual when risk perception itself is low. Therefore, for the purpose of analysing the vaccine intention of individuals, we identify three categories viz., responsive (high risk-high efficacy), denial (high risk-low efficacy) and those with low risk perception (that includes both the unmotivated and indifferent categories). For the group with low risk perception, the model signifies that fear may add to the motivation to process the message cognitively.

### 3. Data

The study uses primarily the fourth wave of the National Income Dynamics Study - Coronavirus Rapid Mobile Survey (NIDS-CRAM 2021) survey. The NIDS-CRAM survey is a special follow up with a subsample of adults from households in Wave 5 of the National Income Dynamics Study (NIDS) run by Southern Africa Labour and Development Research Unit (SALDRU). The NIDS is South Africa's first nationally representative household level panel survey undertaken approximately in two-year intervals between 2008 and 2017, covering a wide range of socio-economic, health, labour and household related information. The NIDS-CRAM survey is designed to be nationally representative and despite the lower sample size (compared to NIDS) remains the best available source of quantitative information on a national scale to assess the socio-economic impact of the corona virus pandemic in South Africa. More detailed technical information on NIDS-CRAM surveys is provided by Ingle et al (2020), Kerr et al (2020), Branson et al (2020) and Daniels et al (2020).

The fourth wave of the NIDS-CRAM has a sample, covering complete questionnaire information, for 5629 individuals. The fourth wave of the NIDS-CRAM survey was conducted over the months of February (96%) and March (4%) 2021. Therefore, the questions were administered under adjusted alert level 3 (Feb 2-28 Feb 2021, when curfew restrictions were imposed and businesses had to adhere to regulations on the maximum number of persons on their premises) and adjusted alert level 1 (March 2021, when most normal activity was allowed to resume with precautions and health guidelines being followed) lockdown conditions.

The wave 4 of NIDS-CRAM introduced a question on vaccine intention “*To what extent do you agree or disagree with the statement: If a vaccine for COVID-19 were available, I would get it?*”, allowing for response on a four-point scale of “strongly agree, somewhat agree, somewhat disagree and strongly disagree”. A further vaccine related question included is “*What is the main reason why you would not take a vaccine for COVID-19?*”.

The key risk perception variable is derived from the NIDS-CRAM questionnaire using the ‘yes’ or ‘no’ response to the question ‘*Do you think you are likely to get the corona virus?*’. The limitation here is that the risk question gives an indication of vulnerability or susceptibility, but not of severity of risk. The study therefore uses two proxy variables to control for the severity of risk based on the emerging evidence that older age groups as well as those with pre-existing chronic illnesses have higher risk of severe COVID symptoms, hospitalization as well as mortality rates compared to younger age groups (Gesese et al 2021, Pijls et al. 2021). Therefore, the first proxy for severity (mortality threat) incorporated in the model is age (measured in years) and the second proxy is obtained by using the ‘yes’ or ‘no’ response to the question “Do you have any of these chronic conditions (you don’t have to tell us which one): HIV, TB, lung condition, heart condition or diabetes?”.

The efficacy variable is sourced from the ‘yes’ or ‘no’ response to the question “*Can you avoid getting Coronavirus?*” The key limitation relating to this variable is that the survey instrument does not afford an opportunity to differentiate between NPI and vaccine related efficacy separately. Given that at the time of the survey vaccines were not yet widely available in South Africa, it is more likely that the responses relate to NPI rather than vaccines. Therefore, we additionally make use of the question “*What is the main reason why you would not take a vaccine for COVID-19?*”, to create a vaccine specific efficacy variable that takes the value one if ineffectiveness of vaccine is not a response to the question. The NPI efficacy and vaccine efficacy variables are interacted to create a combined measure of efficacy.

Lastly, barriers to behavioural change are an important consideration of HBM. One of the barriers is the lack of information. In order to assess the COVID19 related awareness of respondents, a binary variable is constructed which takes the value 1 if the respondent is aware of the three most important symptoms of infection (cough, breathlessness and fever). The assumption made is that a person who is fully aware of the COVID19 symptoms would also be aware of the available information related to vaccine. A further point to note is that the information on knowledge of COVID symptoms was collected in wave 1 during May/June 2020 and as such is limited by the time lapse between the survey waves. More generally, education is also considered as a proxy for awareness.

Apart from awareness, the feasibility of adopting the protective measure is also proxied with household per capita income based on the argument that income reduces the barriers to behavioural change. The household income variable has a high proportion of missing information in the survey data. As such, rand value responses to household income variable have been supplemented with the median value of the income bracket responses. This approach can however distort the income distribution and therefore following Wittenberg (2017) we reweight those who provide rand amounts using the inverse of the probability that an individual will report a rand amount in that bracket. Although this approach does not account for observations that have both the rand value and income bracket information missing, we are able to retain 5272 out of 5629 observations. Further, alternate models are estimated where socio-economic status is controlled for by including other variables such as recent experience of hunger in the household, presence of grant recipients in the household, and electrified dwelling as additional checks.


Because individual responses emerge from a complex interaction of different social, cultural, political and personal factors in vaccine decision (Callaghan et al 2020), additional socio-economic variables such as sex(male), race (black African), partnered, location (urban) and religiosity are incorporated to get a clearer picture of the range of possible predictors about vaccine intention.

Further limitation of the data used is that it is largely self-reported and therefore hard to assert that there was no strategic bias in response. Therefore, the study acknowledges the limitation that it may be susceptible to hypothetical and strategic bias (Althubaiti 2016) especially for key questions like vaccine intention and non-pharmaceutical protective behaviour.

### 4. Preliminary Analysis

Based on the responses to the question on vaccine intention, 55% of the population are seen to have strong acceptance of the vaccine (*Figure 1*). A further 16% are found to accept the vaccine, but with lower certainty. Strong vaccine rejection is indicated by 16%, with a further 7% tending to less strongly reject the vaccine. In addition, 6% did not respond to the question as they were unsure, which may be construed as indicating uncertainty regarding vaccine intention.

**Fig1: If a vaccine for COVID-19 were available, I would get it?**



Source: NIDS-CRAM wave 4. n=5613, weighted.

Converting the four-scale response to binary variable by combining 'strongly agree' with 'somewhat agree' to define vaccine acceptance; and 'strongly disagree' with 'somewhat disagree' as well as 'don't know', to derive the vaccine hesitancy. Based on this definition, vaccine hesitancy is calculated as being 29.16% (with 95% confidence interval of 28%-30%). This is a significant proportion of population and raises concern regarding the success of achieving herd immunity through vaccination.

**Table 2: Summary statistics of Risk and Efficacy, by severity/mortality risk**

Variable	Mean	Age: 60 years and above	Age: below 60 years	t-test	chronic	no chronic	t-test
Risk	0.450 (0.498)	0.398 (0.490)	0.458 (0.498)	*	0.481 (0.500)	0.444 (0.497)	-
Efficacy	0.824 (0.381)	0.829 (0.376)	0.823 (0.382)	-	0.850 (0.357)	0.818 (0.386)	-
High Risk, High efficacy	0.305 (0.460)	0.216 (0.412)	0.321 (0.467)	***	0.339 (0.474)	0.297 (0.457)	-
High Risk, Low efficacy	0.064 (0.245)	0.032 (0.176)	0.070 (0.255)	***	0.055 (0.227)	0.066 (0.248)	-
Low -risk	0.550 (0.498)	0.602 (0.490)	0.542 (0.498)	*	0.519 (0.500)	0.556 (0.497)	-
Low risk, Low efficacy	0.080 (0.077)	0.077 (0.266)	0.081 (0.273)	-	0.070 (0.255)	0.082 (0.275)	-
Low risk, High efficacy	0.376 (0.484)	0.343 (0.475)	0.382 (0.486)	-	0.346 (0.476)	0.382 (0.486)	-

**Source:** NIDS-CRAM wave 4, weighted. Standard errors in parentheses, \*\*\* p<0.01, \*\* p<0.05, \* p<0.1


Considering the risk and efficacy variables, it is clear that the risk perception (45%) is substantially lower than efficacy (82%) (Table 2). While 45% perceived the risk of COVID19 infection, a higher proportion of 55% did not. Age and co-morbidities are considered as two risk factors that increase the probability of severe symptoms, hospitalization and mortality. As such we examine whether risk and efficacy variables are different for those in these categories. The average risk perception is higher for those below sixty years of age, explained by the elderly more likely to be staying home with lower external exposure. The corollary of this is that higher proportion of elderly perceive low risk compared to the younger group. A significantly lower proportion of the elderly are in the responsible and denial categories compared to those below sixty years. Within the EPPM risk-efficacy categories, the highest proportion is of low risk-low efficacy (indifference) group (48.3%). Responsive group is estimated at 39.5% of the adult population. No significant difference is found in the risk and efficacy levels of those suffering from chronic conditions and others.

## 5. Assessing vaccine hesitancy

Figure 2 presents average vaccine hesitancy levels across sex, race and education attainment. While females in the sample had on average higher vaccine hesitancy (31%) compared to males (27%), the difference is significant at 95% confidence level. What is clearly evident at even 99% confidence level is the higher vaccine hesitancy among non-black population (35%), compared to the black African population (23%). Those with tertiary education is seen to have significantly lower hesitancy (26%) compared to those with below tertiary education (31%). Geographical location is found to be also a significant differentiating factor with vaccine hesitancy found to be higher in urban areas (30%) compared to rural areas (27%). Individuals who reported chronic illness have on an average lower (23%) vaccine hesitancy compared to those who did not report chronic illness (30%).


Considering the association between health behaviour variables and vaccine hesitancy, risk perception is seen to be more significant than efficacy as a differentiating variable of vaccine hesitancy (Figure 3). Individuals with high-risk perception have statistically lower vaccine hesitancy (26%) compared to those with low-risk perception (31%). Statistically significant differences are evident in the hesitancy across the efficacy groups as well as the risk-efficacy interaction groups (responsive and denial).

**Fig 2 Vaccine hesitancy, Sex, Race and Education**




Source: NIDS-CRAM wave 4, weighted.

**Fig 3 Vaccine hesitancy, Risk and Efficacy**



Source: NIDS-CRAM wave 4, weighted.

**Fig: 4 Vaccine hesitancy and key characteristics**



Source: NIDS-CRAM wave 4, weighted.

Figure 4 highlights significant difference between the vaccine hesitancy rates of urban and rural areas, those who reported chronic illness and those who did not, those who had full information and those who did not, who experienced hunger and who did not. While these findings are relevant and points to possible predictors, it is possible that there may be confounding factor like income behind some of these variables. For instance, poverty and hunger is higher in rural areas in South Africa. Thus, it is important to undertake a multivariate analysis before drawing firm conclusions on the predictors of vaccine hesitancy.

## 6. Socio-economic inequality in vaccine hesitancy, risk and efficacy

This section uses concentration index to assess the socio-economic inequality in the prevalence of vaccine hesitancy, as well as the risk and efficacy parameters along the income, age and education distributions. As vaccine hesitancy, risk perception and efficacy variables are all binary, Erreygers-corrected concentration index (Erreygers 2009) is used to assess their socio-economic inequality.

While income-based inequality is not evident for vaccine hesitancy, it is clear that income plays a key role when it comes to risk and efficacy (*Table 3*). Vaccine hesitancy is negative and significant along age, pointing to higher acceptance among the elderly. The higher hesitancy concentration among the young could be related to the perceived lower severity of the infection related morbidity and mortality among the younger age groups (De Bruin 2021, Nino et al 2021).

Risk perception is significantly higher among those with higher income. Similar results have been reported by Kollampambil and Oyenubi (2021) based on wave 1 and 2 of the NIDS-CRAM. Our results show that this pattern is sustained after a year. As expected, risk perception has a significant positive concentration along the age distribution indicating higher risk perception among the older individuals (*Table 3*). This is not surprising considering the available information that age increases severity of COVID illness, the risk of hospitalisation and higher COVID infection related mortality rate (Gesese et al 2021, Pijls et al 2021). This is related to the results observed with regards to vaccine hesitancy, and points to the mediating role of risk perception in leading to the higher vaccine hesitancy among younger age groups.

Both risk perception and efficacy are found to be pro-education. The risk-efficacy interaction variables are led by risk in terms of their concentration patterns. 'Responsive' (high risk-high efficacy) and 'Denial' (high risk-low efficacy) are both pro-rich, aligned with the results of risk perception. The low risk category (combination of low-risk-high efficacy, and low risk-low efficacy groups) are pro-poor. Low-risk are also pro-young and pro-low education.

The findings indicate that although vaccine hesitancy is not concentrated along the income or education distribution, they can nevertheless play a role through mediating factors like risk perception which have definite concentration along income, age and education distributions.

**Table 3 Erreygers concentration index of vaccine hesitancy, risk and efficacy**

CI	Vaccine hesitancy	Risk	Efficacy	Responsive (High risk-high efficacy)	Denial (High risk-Low efficacy)	Low risk
Income	-0.017 (0.031)	0.223*** (0.035)	-0.03 (0.024)	0.143*** (0.030)	0.043*** (0.016)	-0.223*** (0.35)
Age	-0.121*** (0.023)	0.111*** (0.026)	-0.003 (0.017)	0.023 (0.023)	-0.014 (0.009)	-0.111*** (0.026)
Education	-0.018 (0.023)	0.132*** (0.027)	0.045*** (0.018)	0.114*** (0.025)	0.029** (0.011)	-0.133*** (0.027)

Standard errors in parentheses, \*\*\* p<0.01, \*\* p<0.05, \* p<0.1

## 7. Multivariate analysis

The descriptive statistics of the key socio-economic characteristics of the sample for the multivariate regression indicate resonance with national statistics (*Table 4*).

**Table 4: Summary statistics of the Study Sample**

Variable	Obs	Mean	Std. Dev.	Min	Max
Age, years (Risk Severity risk 1)	4,440	40.47	15.04	18	90
Chronic illness (Risk Severity 2)	4,440	0.168	0.374	0	1
Well-informed (Barrier 1)	4,440	0.096	0.295	0	1
Household per capita income, Rands (Barrier 2)	4,440	2854	6135	0	150000
Education, years	4,440	11.52	3.729	0	22
African	4,440	0.782	0.413	0	1
Male	4,440	0.482	0.500	0	1
Married/with partner	4,440	0.479	0.500	0	1
Urban	4,440	0.760	0.427	0	1
Informal_dwelling	4,324	0.117	0.322	0	1
Electricity	4,440	0.946	0.226	0	1
Experienced Hunger	4,440	0.164	0.370	0	1
Employed	4,440	0.497	0.500	0	1
Social Grant recipient	4,440	0.353	0.478	0	1
Religious	4,440	0.913	0.282	0	1

Source: NIDS-CRAM weighted. # All variables except age, education and income are binary

The results of logit regressions highlight the role of risk and efficacy in driving vaccine hesitancy while controlling for other pertinent predictors (*Table 5*). Columns 1 - 3 and 4-6 are based on the HBM, and the EPPM respectively. Various specifications are estimated using log of per capital income, income quintiles and non-financial proxies for socio-economic status (hunger in household, household with electricity, grant recipient in household).

**Table 5: Logit regression results: Dependent variable, Vaccine Hesitancy**

VARIABLES	(1)	(2)	(3)	(4)	(5)	(6)
	HBM	HBM	HBM	EPPM	EPPM	EPPM
Risk perception	-0.265** (0.133)	-0.268** (0.133)	-0.271** (0.133)			
Efficacy	-1.260*** (0.155)	-1.260*** (0.154)	-1.248*** (0.153)			
Low risk^				0.429*** (0.125)	0.433*** (0.124)	0.429*** (0.124)
Denial^				1.061*** (0.230)	1.068*** (0.230)	1.052*** (0.231)
Severity 1(Age)	-0.0251*** (0.00539)	-0.0260*** (0.00531)	-0.0262*** (0.00534)	-0.0216*** (0.00472)	-0.0220*** (0.00469)	-0.0221*** (0.00469)
Severity 2 (chronic)	-0.155 (0.175)	-0.153 (0.175)	-0.158 (0.176)	-0.263* (0.155)	-0.267* (0.155)	-0.241 (0.156)
Barrier 1 (Information)	-0.287 (0.217)	-0.330 (0.215)	-0.338 (0.218)	-0.431** (0.213)	-0.459** (0.212)	-0.469** (0.213)
Barrier 2 (education)	-0.0442** (0.0191)	-0.0511*** (0.0196)	-0.0499*** (0.0190)	-0.0370** (0.0177)	-0.0414** (0.0184)	-0.0437** (0.0176)
Barrier 3 (log PC HH Income)	-0.071* (0.0418)			-0.0694* (0.0389)		
2. PC HH Income _Q2		0.036 (0.177)			0.068 (0.28)	
3. PC HH Income _Q3		-0.348 (0.223)			-0.271 (0.194)	
4. PC HH Income _Q4		-0.101 (0.201)			-0.0545 (0.178)	
5. PC HH Income _Q5		0.0524 (0.204)			0.0118 (0.191)	
Hunger			-0.216 (0.167)			-0.212 (0.149)
Electricity			0.135 (0.302)			-0.109 (0.259)
Grant receiving Household			0.210 (0.132)	0.136 (0.117)	0.144 (0.118)	0.154 (0.118)

African	-0.817*** (0.216)	-0.768*** (0.214)	-0.782*** (0.215)	-0.776*** (0.196)	-0.743*** (0.194)	-0.718*** (0.194)
NPI behaviour	-0.114** (0.0578)	-0.117** (0.0577)	-0.119** (0.0577)	-0.140*** (0.0512)	-0.142*** (0.0513)	-0.148*** (0.0511)
Male	-0.218* (0.125)	-0.240* (0.126)	-0.216* (0.124)	-0.169 (0.114)	-0.182 (0.115)	-0.174 (0.113)
Married/Partnered	0.214 (0.132)	0.205 (0.131)	0.228* (0.133)	0.251** (0.120)	0.250** (0.119)	0.237** (0.120)
Employed	0.196 (0.130)	0.165 (0.133)	0.178 (0.131)	0.0406 (0.121)	0.0202 (0.122)	-0.0193 (0.120)
Religious	-0.0867 (0.240)	-0.0657 (0.240)	-0.0872 (0.239)	0.00923 (0.223)	0.0305 (0.226)	0.0226 (0.222)
Province controls	yes	yes	yes	yes	yes	yes
Constant	3.037*** (0.635)	2.529*** (0.579)	2.420*** (0.676)	1.544** (0.601)	1.021* (0.556)	1.235** (0.613)
<b>Wald chi2</b>	<b>138.57**</b>	<b>158.31***</b>	<b>158.14***</b>	<b>115.46***</b>	<b>122.23***</b>	<b>115.96***</b>
<b>Observations</b>	<b>4,264</b>	<b>4,264</b>	<b>4,235</b>	<b>4,441</b>	<b>4,441</b>	<b>4,440</b>

**Robust** standard errors in parentheses, \*\*\* p<0.01, \*\* p<0.05, \* p<0.1  
 ^ the benchmark variable is Responsive (High risk-high efficacy)

The results of HBM specifications are consistent in underscoring the significance of risk perception of infection and efficacy as negative predictors of vaccine hesitancy. The EPPM results also indicate that individuals who perceive lower vulnerability to infection are more likely to be hesitant about accepting vaccine compared to the responsive (high risk and high efficacy) group.

The results indicate that, compared to responsive individuals, the low-risk individuals are more likely to be hesitant regarding the vaccine. We also see a significant difference between the vaccine intention of the 'responsive' and the 'denial' groups. This once again points to risk perception as being the driving factor, with no clear mediating role of efficacy being apparent.

Therefore, EPPM estimations further highlight the mediating role of efficacy, where the denial group (high risk-low efficacy) has a significantly higher vaccine hesitancy in relation to the responsive group. These results are insightful and suggest that it is important to accompany risk messaging with efficacy related communication to encourage individuals to accept vaccination.

Severity of risk, proxied through age has a significant negative association with vaccine hesitancy in both HBM and EPPM estimations. The second proxy for severity, chronic illness, is also significant in the EPPM model. The indication therefore is that in addition to vulnerability to infection, vaccine intentions are strongly driven by the perceived severity of risk as well.

Income, education and information availability are incorporated in the model as measures of barriers to adoption of health response. Education is likely to be correlated with both income and Covid related information, as such its significance is indicative of the key role that awareness plays in vaccine intentions. Similarly, the variable relating to information regarding COVID is negative (although significant only in the EPPM estimations) pointing to the need for creating more awareness and education around vaccines to reduce hesitancy. The income variable is rather weak and does

not come out strong in the estimations. This however could be due to its strong correlation with some of the other variables in the model like education. The other proxies included to control for socio-economic status like experience of hunger, social grant receipt, and electrified dwelling also do not give any strong indications on their role in driving vaccine hesitancy. However, there is still the possibility that socio-economic status acts via risk perception and efficacy. This is explored further in section 10.

The positive association between adoption of NPI behaviour (measured as number of non-pharmaceutical preventive behaviour adopted) and vaccination is evident in both the HBM and EPPM models. There are other interesting socio-economic predictors of vaccine intention revealed through both HBM and EPPM estimations. Black Africans have significantly lower vaccine hesitancy compared to other race groups. Being married or having a partner is found to increase vaccine hesitancy significantly. This finding is contrary to expectations as being married is found in literature to have protective effect on health behaviour (Umberson 1992). However, there have been studies (Thompson et al 2017) that cite negative relationship between vaccination and being married (or in a steady relationship). The result that males are on average less hesitant than females is consistent with the finding by Edwards et al (2021) in the Australian context.

In order to remove any bias caused by the construct of the efficacy variable, we re-estimate the HBM and EPPM specifications excluding the observations where only vaccine hesitancy drives the efficacy variable. In other words, we drop observations where vaccine ineffectiveness takes the value one and the general efficacy variable is one. This ensures that there is no definitional relationship between the efficacy variable and the dependent variable of vaccine hesitancy. The results from this restricted estimation are given in appendix (*Table A1*) and validates the results in *Table 5*.

As a further robustness check we also present results of ordinal logit regressions, where the dependent variable of vaccine intention is used in its original form with four response categories. The marginal effects of the regression relating to the key risk and efficacy variables for each of the four outcomes are presented in *Table 6*.

The results are consistent with the logit regression with regards to risk and efficacy for the HBM model, and the risk-efficacy interaction variables (denial and low risk) for the EPPM model. The findings hold that risk perception, through the mediation of efficacy is driving vaccine intentions. The severity of risk proxied by age is significant predictor as in the logit model. There are however some deviations from the logit model in relation to chronic illness with it being significant in HBM but not in EPPM estimation. Education on the other hand, is significant only in the EPPM model for the ordinal estimation.

**Table 6 Ordinal logit, marginal effects: Dependent variable, Vaccine Intention.**

HBM	High Risk	High Efficacy	Severity-age	Severity-Chronic illness	Barrier-Education	Barrier-income	Barrier-information
Strongly accept vaccine	0.050** (0.024)	0.197*** (0.031)	0.007*** (0.001)	0.070** (0.032)	0.003 (0.003)	0.016 (0.011)	0.047 (0.045)
Somewhat accept vaccine	-0.012** (0.005)	-0.048*** (0.008)	-0.002*** (0.000)	-0.017** (0.008)	-0.001 (0.001)	-0.004 (0.003)	-0.011 (0.011)
Somewhat reject vaccine	-0.009** (0.004)	-0.038*** (0.009)	-0.001*** (0.000)	-0.014** (0.006)	-0.001 (0.001)	-0.003 (0.002)	-0.010 (0.009)
Strongly reject vaccine	-0.028** (0.016)	-0.111*** (0.024)	-0.004*** (0.001)	-0.039** (0.020)	-0.002 (0.002)	-0.009 (0.006)	-0.026 (0.025)

EPPM	Denial ^	Low risk^	Severity-age	Severity-Chronic illness	Barrier-Education	Barrier-income	Barrier-information
Strongly accept vaccine	-0.173*** (0.044)	-0.066*** (0.024)	0.007*** (0.001)	0.055 (0.036)	0.008** (0.002)	0.016 (0.011)	0.039 (0.044)
Somewhat accept vaccine	0.040*** (0.011)	0.016*** (0.005)	-0.002*** (0.000)	-0.013 (0.009)	-0.002** (0.000)	-0.004 (0.003)	-0.010 (0.011)
Somewhat reject vaccine	0.035*** (0.003)	0.013*** (0.006)	-0.002*** (0.000)	-0.012 (0.008)	-0.002** (0.000)	-0.003 (0.002)	-0.008 (0.009)
Strongly reject vaccine	0.037*** (0.018)	0.098*** (0.025)	-0.004*** (0.001)	-0.030 (0.020)	-0.005** (0.000)	-0.009 (0.006)	-0.022 (0.024)

**Robust** standard errors in parentheses, \*\*\* p<0.01, \*\* p<0.05, \* p<0.1, ^Benchmark category is responsive (high risk-high efficacy)

## 8. Reasons for Vaccine hesitancy

In this section we look at reasons reported by respondents for vaccine hesitancy. The analysis is based on the response to the question in the survey “*What is the main reason why you would not take a vaccine for COVID-19?*”. Among the reasons cited for vaccine hesitancy, the highest proportion cited concerns regarding side-effects of vaccines (30%) (Figure 5). This is not surprising considering the lack of public information on vaccine trials as well as emerging media reports on possible serious side-effects of certain vaccines like blood-clotting (CIDRAP 2021).


Concerns regarding the side-effects of vaccine aside, almost 19% of those who indicated unwillingness to accept vaccine pointed to ineffectiveness of vaccines as a reason for vaccine hesitancy. The mixed messaging coming through media on the efficacy of vaccines as well as their uncertainty regarding the usefulness against the new COVID strains are contributing to the concerns regarding ineffectiveness of vaccine. The concerns regarding side effects and ineffectiveness (jointly accounting for 49%) are specific to COVID19 vaccines and therefore needs to be addressed with specific information on COVID19 vaccines. A further 2% indicated lack of trust in vaccines and government. The lack of clarity in communication and unavailability of information on vaccines are clearly contributing to vaccine hesitancy.

General anti-vaccine stance, that extends beyond COVID specific vaccine inhibition accounted for nearly 15% of individuals who indicated vaccine hesitancy. This category of individuals will need to be addressed with information beyond COVID-specific vaccine. The approach taken would have to be broader, but nevertheless COVID specific information will also be useful in generating awareness on vaccine efficacy.

The response efficacy related factors aside, it is clear from that a significant section of individuals do not believe that COVID poses a significant risk (10%). The need to address this category with risk messaging is clearly evident. Further a prominent reason cited within the “other” category is “conflicting messages from media / on social media”.

The reasons for vaccine hesitancy therefore highlight the need to improve communication regarding efficacy of vaccines as well as to create further awareness on the risks associated with COVID through risk messaging.

**Figure 5: Reasons for Vaccine hesitancy (%)**



Source: NIDS-CRAM wave 4, weighted.

**Table 7: Erreygers-corrected Concentration index for vaccine hesitancy reasons**

CI	Side-effects	Anti-vaccination	Ineffective	No COVID risk
income	-0.069	-0.006	0.117**	0.016
SE	(0.063)	(0.056)	(0.051)	(0.034)
age	0.050	-0.035	-0.138**	0.030
SE	(0.053)	(0.043)	(0.049)	(0.024)
education	0.010	-0.062	0.101*	-0.024
SE	(0.048)	(0.045)	(0.046)	(0.029)

Robust standard errors in parentheses, \*\*\* p<0.01, \*\* p<0.05, \* p<0.1

There is not much indication of concentration of the various reasons for vaccine hesitancy along the income, age and education distribution, with the exception of ‘ineffective’ as a reason (Table 7). ‘Ineffective’ is found to be pro-young and pro-educated. The scepticism regarding the efficacy of vaccination needs to be communicated within these socio-economic groups through more transparent sharing of information based on scientific research.

## 9. Comparison with Non-Pharmaceutical intervention (NPI)

Studies on the adoption of NPI in South Africa indicate a high level of uptake, with close to 100% of the population indicating some form of changed behaviour as a protective response to COVID. This is much higher than the vaccine acceptance currently observed. It is not surprising that inhibitions regarding vaccines are higher than NPI, and can be attributed to possible secondary harms through vaccination. Also, unlike vaccination, some NPI measures like wearing of facemask and limits on social gatherings have been made mandatory via government regulation. This implies that adoption of these mandatory NPI measures may be less driven by individual perception of risk and efficacy.

The results of the seemingly unrelated regressions (*Tables A2 and A3*) validate this where the role of efficacy is less evident for NPI for the most popular non-pharmaceutical preventive behaviour (the use of facemask) in South Africa (Kollamparambil & Oyenubi, 2021). Efficacy however has a significant association with the use of sanitizers and hand hygiene behaviour. Similarly, the mediating role of efficacy is not strongly evident for facemask use, but is significant for the practise of hand hygiene and the use of sanitizers. The indications are that the government regulation making facemask use mandatory has impacted on behaviour and therefore it is less driven by individual discretion which is influenced by efficacy.

The role of risk perception on the other hand, is found to have a significant association across all forms of non-pharmaceutical protective behaviour as well as vaccine intention (*Tables A2, A3 and Table 5*). The low-risk group has significantly lower probability of adopting the most popular forms of NPI indicating that NPI behaviour is driven primarily through risk assessment.

## 10. Targeted communication

The multivariate analysis of predictors of vaccine hesitancy has identified denial and low-risk perception (low risk-low efficacy and low risk- high efficacy) categories as positive predictors of vaccine hesitancy. Therefore, it is important to target foremost these groups for health communication based on not just fear appeal but also efficacy messaging. This requires identifying the socio economic and demographic characteristics of these groups. For this purpose, the study next undertakes a multinomial logit regression, where the base category is the responsive (high risk-high efficacy) category (*Table 8*). It must be flagged here that the construct of the risk categories here are based on risk perception and not severity of risk.

**Table 8: Multinomial logit regression: Base regression, Responsive category.**

VARIABLES	Denial <sup>^</sup>	Low risk <sup>^</sup>
Age	-0.0152 (0.0419)	-0.105*** (0.0227)
Age squared	1.77e-05 (0.000428)	0.00109*** (0.000249)
Education (years)	0.00383 (0.0376)	-0.0445** (0.0175)
African	-0.637* (0.368)	0.107 (0.222)
Well informed	0.0925 (0.312)	-0.290 (0.192)
Male	0.313 (0.225)	-0.00667 (0.118)
Married	-0.104 (0.229)	-0.304** (0.122)
Urban	0.190 (0.323)	-0.0475 (0.126)
Informal_dwelling	-0.0408 (0.410)	-0.260 (0.196)
Electricity	0.00573 (0.711)	-0.474* (0.262)
Employed	-0.203 (0.267)	-0.665*** (0.132)
Household income	-0.0104 (0.0839)	-0.103** (0.0438)
Grant recipient	0.283 (0.256)	-0.145 (0.121)
Religious	1.157** (0.529)	-0.165 (0.228)
Chronic illness	-0.210 (0.270)	-0.242 (0.148)
Province control	yes	yes
Constant	-2.328 (1.839)	4.920*** (0.735)
<b>Wald chi2(46)</b>	<b>206.46***</b>	<b>206.46***</b>
<b>Observations</b>	<b>4,179</b>	<b>4,179</b>

**Robust** standard errors in parentheses, \*\*\* p<0.01, \*\* p<0.05, \* p<0.1

<sup>^</sup> Base regression: Responsive (High risk-high efficacy)

Compared to the benchmark category (responsive), higher age is likely to decrease the probability of belonging in the low-risk group (*Table 7*). However, the non-linear relationship indicates that the elderly is also likely to belong to the low-risk group, explained due to the fact that they are mostly home-bound and not exposed to external interactions. Education is also likely to decrease the probability of belonging in the low-risk perception group, vis-à-vis the responsive group. Being married (or with a partner) as well as being employed has a similar effect of reducing the probability of being in the low-risk perception group. Higher socio-economic status (as indicated by income, being employed and electrified dwelling) also decreases the probability of being in the low-risk perception group.

There are not many significant identifiers of the denial group (high risk-low efficacy) from the responsive group (high risk-high efficacy). While black African have lower probability of being in the denial category compared to responsive category, those who consider religion to be important are more likely to be in the denial category.

These results indicate that targeting the young, the less informed, low-income individuals, those without partners, and the unemployed through risk messaging. On the other hand, non-blacks and the religious need to be targeted for protective danger control and education on solutions to the threat of the pandemic. Over 91 percent report that religion plays an important role in their life, and as such our results show that efficacy communication through religious leaders and institutions would be effective.

Our findings show that unless efficacy of the available response behaviour (NPI and vaccine) is communicated, enhanced risk perception alone will not result in appropriate behavioural response (including vaccine acceptance).

## 11. Summary and Policy implications

Based on the vaccine intentions of the respondents of NIDS-CRAM survey, only 55% are fully accepting of the vaccine, while a further 16% are moderately accepting of vaccines. Combining these two groups, vaccine acceptance is estimated at 70.8% in South Africa. Correspondingly, vaccine hesitancy in South Africa is estimated at 29.2% (with 95% confidence interval of 28%-30%) for the general adult public. This is significantly high and raises concern regarding the effectiveness of vaccination programme once vaccines become widely available to the public. This is especially when considering that vaccines are not 100% effective, and for vaccine effectiveness of 60%, vaccine coverage of 100% is required to achieve control of the virus (Bartsch et al 2020). These results relate to information collected in February 2021 and needs to be monitored closely going forward.

Comparing the findings of this study with the meta-analysis by Sallam (2021), vaccine hesitancy in South Africa is higher than countries like Ecuador, Malaysia, Indonesia and China; but lower than countries like Kuwait, Jordan, Italy, Russia, Poland, US, and France.

Vaccine acceptance in South Africa is substantially lower than the non-pharmaceutical behavioural changes undertaken as protective measure against COVID19. Despite the sustained efforts required for NPI based preventive strategy, the concerns regarding the perceived dangers of secondary harm of pharmaceutical interventions compared to NPIs is an emerging explanation for this. Another possible explanation is that some non-pharmaceutical measures like the use of face mask and limits to social gathering have been made mandatory under government regulations, whereas vaccine acceptance is considered to be left to individual discretion. Nonetheless, the results indicate that individuals who take NPI seriously are also more likely to accept vaccination.

There are significant differences across socio-economic variables in vaccine hesitancy. Age has emerged as a significant predictor of vaccine hesitancy with younger individuals more likely to be against vaccine acceptance. This is not surprising given the lower risk perception among the young

with regards to the severity of COVID19. Similarly, those reporting chronic illness also have lower hesitancy compared to their healthier counterparts.

Non-black African population group has significantly high vaccine hesitancy compared to black Africans. Males on average have lower hesitancy compared to females. Unmarried/unpartnered individuals are found to have higher hesitancy. Education on the other hand is found to reduce hesitancy.

In addition, health behaviour models offer important predictors of vaccine intentions. Risk perception (both vulnerability and severity) is a key driving factor along with efficacy. This study validates the argument by Witte (2002, 2004) that efficacy plays an important mediating role in the relationship between risk perception and the associated behavioural response. The group that was seen to be most accepting of vaccines are those that perceive high risk as well as high efficacy. Doubts regarding the response efficacy of vaccination however has emerged as the deterring reason against vaccine acceptance. From a communication perspective, it is imperative to continue risk messaging, hand in hand with clearer information on the efficacy of the vaccines.

The fear messaging needs to be targeted at younger age population who perceive the lowest risk. Those who are less informed on COVID19 are also found to be those who perceive low risk; therefore, this reinforces the need to have in place a communication plan to educate and create awareness on the pandemic. However, as highlighted by Oyenubi et al (2021) excessive fear messaging without efficacy-related messaging can lead to mental health issues. Therefore, efficacy messaging cannot be ignored. Considering that 91% of South Africans consider religion to be important, and the finding that religious individuals are more likely to belong to the denial (high risk-low efficacy) group, a recommendation emerging is the use of religious leaders and institutions for efficacy messaging.

The need for efficacy messaging also is showcased by an analysis of the reasons that were cited for vaccine hesitation. These include primarily the possible negative side-effects, and the ineffectiveness of vaccine. While the risk of blood clots associated in particular with AstraZeneca and Johnson & Johnson vaccines has grabbed headlines, the risk of the vaccine vis a vis more accepted pharmaceutical practices like the use of contraceptive pills is considered to be lower (Taylor 2021). This shows that there is a need to make available more information from the scientific studies on the possible side-effects and effectiveness that would allow individuals to make more informed judgements. The lack of transparency in this regard is contributing substantially to vaccine hesitancy and there is a clear need for better communication around this.

General anti-vaccine stance, that extends beyond COVID specific vaccine inhibition accounted for nearly 15% of individuals who indicated vaccine hesitancy. This category of individuals will need to be addressed with information beyond COVID-specific vaccine. The approach taken would have to be broader, but nevertheless COVID specific information will also be useful in generating awareness on vaccine efficacy.

**Table 9: Communication strategy recommendations**

---

<p>Responsive: People are educated about risk, believe in efficacy and respond by taking protective action. 40% of population are responsive. Vaccine hesitancy is 26%. Profile: Older, educated, married, well-informed Strategy: Provide calls to vaccination and make vaccination available. Provide clear information on COVID19 vaccine side effects and effectiveness</p>	<p>Denial: People are aware of the risks but feel helpless to respond effectively. 7% of population fall in this category. Vaccine hesitancy is 27%. Profile: Non-black, religious Strategy: Less focus on fear messaging and more on protective danger control through educating about solutions Provide clear information on COVID19 vaccine side effects and effectiveness</p>
<hr/>	
<p>Low risk: Due to the low risk perception, people do not enter into efficacy assessment. 52.6% of population fall in this category Vaccine hesitancy is highest at 31%. Profile: Young, low education, single status, less informed, low income Strategy: Educate through risk and fear messaging, followed by risk and protective danger control through awareness about solutions. Provide clear information on COVID19 vaccine side effects and effectiveness</p>	

---

Widening vaccine acceptance requires a three-pronged health communication strategy being put in place that targets fear messaging to the young, uneducated, less informed, lower income individuals; efficacy messaging to non-black and religious individuals; call to action among older, educated and well-informed individuals. The targeted communication aside summarized in *Table 9*, there is an over-arching need to provide research-based information on the side-effects and effectiveness of COVID 19 vaccines across the board.

## REFERENCES

- Althubaiti A. 2016. Information bias in health research: definition, pitfalls, and adjustment methods. *Journal of multidisciplinary healthcare*, 9: 211. doi:10.2147/JMDH.S104807.
- Barnett DJ, Errett NA, Rutkow L. 2013. A Threat- and Efficacy-Based Framework to Understand Confidence in Vaccines among the Public Health Workforce, *Vaccines*, 1, 77-87; doi:10.3390/vaccines1020077
- Bartsch S. M., O'Shea K. J., Ferguson M. C., Bottazi M. E., Wedlock P. T., Strych U., et al. 2020. Vaccine efficacy needed for a COVID-19 Coronavirus vaccine to prevent or stop an epidemic as the sole intervention. *American Journal of Preventive Medicine*, 59(4), 493–503. <https://doi.org/10.1016/j.amepre.2020.06.011> PMID: 32778354
- Boster, F. J., & Mongeau, P. 1984. Fear-arousing persuasive messages. *Annals of the International Communication Association*, 8(1), 330-375.
- Branson N, Wittenberg M. 2019. Longitudinal and cross-sectional weights in the NIDS data 1-5 [NIDS Technical Paper Number 9]. Southern Africa Labour and Development Research Unit, University of Cape Town.
- Burger, R., Christian, C., Maughan-Brown, B., Rensburg, R & Rossouw, L. 2020. COVID-19 subjective risk perception knowledge and behaviour. Wave 1, Report No. 2, NIDS-CRAM.
- Callaghan T, Moghtaderi A, Lueck JA, Hotez P, Strych U, Dor A, Fowler EF, Motta M. Correlates and disparities of intention to vaccinate against COVID-19.2021. *Soc Sci Med.*, Mar;272:113638. doi: 10.1016/j.socscimed.2020.113638. Epub 2021 Jan 4. PMID: 33414032; PMCID: PMC7834845.
- Carcioppolo, N., Jensen, J. D., Wilson, S. R., Collins, W. B., Carrion, M., & Linnemeier, G. 2013. Examining HPV threat-to-efficacy ratios in the extended parallel process model. *Health Communication*, 28(1), 20-28.
- Chu, H and Liu, S. 2021. Integrating health behavior theories to predict American's intention to receive a COVID-19 vaccine, *Patient Education and Counseling*, <https://doi.org/10.1016/j.pec.2021.02.031>.
- CIDRAP.2021. Groups find possible link between AstraZeneca COVID vaccine, blood clots, Centre for Infectious Disease research and Policy, University of Minnesota. Groups find possible link between AstraZeneca COVID vaccine, blood clots | CIDRAP (umn.edu)
- Daniels RC, Ingle K, Brophy T.2020. Determinants of attrition in NIDS-CRAM Waves 1 & 2. NIDS-CRAM.
- De Bruin, W. B. 2021. Age Differences in COVID-19 Risk Perceptions and Mental Health: Evidence From a National U.S. Survey Conducted in March 2020. *The Journals of Gerontology: Series B*, Volume 76, Issue 2, February 2021, Pages e24–e29, <https://doi.org/10.1093/geronb/gbaa074>
- Dillard, J. P. 1994. Rethinking the study of fear appeals: An emotional perspective. *Communication Theory*, 4(4), 295-323.
- Dror, A.A., Eisenbach, N., Taiber, S. et al. 2020. Vaccine hesitancy: the next challenge in the fight against COVID-19. *Eur J Epidemiol* 35, 775–779. <https://doi.org/10.1007/s10654-020-00671-y>
- Dubé E, Laberge C, Guay M, Bramadat P, Roy R, Bettinger J. 2013. Vaccine hesitancy: an overview, *Human Vaccines & Immunotherapeutics*, 9:8, 1763-1773.
- Edwards B, Biddle N, Gray M, Sollis K. 2021. COVID-19 vaccine hesitancy and resistance: Correlates

in a nationally representative longitudinal survey of the Australian population. *PLoS ONE* 16(3): e0248892. <https://doi.org/10.1371/journal.pone.0248892>

Erreygers G. 2009. Correcting the concentration index. *Journal of health economics*;28: 504–515.

Gesesew HA, Koye DN, Fetene DM, et al. 2021. Risk factors for COVID-19 infection, disease severity and related deaths in Africa: a systematic review *BMJ Open* 2021;**11**:e044618. doi: 10.1136/bmjopen-2020-044618

Grundey, S.M. 1999. Age as a risk factor: You are as old as your arteries. *American Journal of Cardiology*, 83, 1455-1457.

Ingle K, Brophy T, Daniels RC. 2020. National Income Dynamics Study–Coronavirus Rapid Mobile Survey (NIDS-CRAM) panel user manual. Technical Note Version. 2020;1. NIDS-CRAM.

Kerr A, Ardington C, Burger R. 2020. Sample design and weighting in the NIDS-CRAM survey. NIDS-CRAM.

Kollamparambil, U. & Oyenubi, A. 2020a. Socio-economic inequality in the response to COVID19 pandemic, Policy paper, wave 2 no. 10, NIDS-CRAM.

Kollamparambil, U. & Oyenubi, A. 2021. Behavioural response to the Covid-19 pandemic in South Africa, *PLoS One*, forthcoming. <https://doi.org/10.1371/journal.pone.0250269>

Latimer, A. E., Brawley, L. R., & Bassett, R. L. 2010. A systematic review of three approaches for constructing physical activity messages: what messages work and what improvements are needed? *International Journal of Behavioral Nutrition and Physical Activity*, 7(1), 36.

Leventhal, H. 1970. Findings and theory in the study of fear communications. In L. Berkowitz (Ed.), *Advances in experimental social psychology* (Vol. 5, pp. 119-186). New York: Academic Press. doi:10.1016/S0065-2601(08)60091-X

Lin Y, Hu Z, Zhao Q, Alias H, Danaee M, Wong LP. 2020. Understanding COVID-19 vaccine demand and hesitancy: A nationwide online survey in China. *PLoS Negl Trop Dis* 14(12): e0008961. <https://doi.org/10.1371/journal.pntd.0008961>

Lin, H. C., & Chen, C. C. 2021. Disease Prevention Behavior During the COVID-19 Pandemic and the Role of Self-Esteem: An Extended Parallel Process Model. *Psychology research and behavior management*, 14, 123–135. <https://doi.org/10.2147/PRBM.S291300>

Lithopoulos, A., Liu, S., Zhang, C.-Q., & Rhodes, R. E. 2021. Predicting Physical Distancing in the Context of COVID-19: A Test of the Extended Parallel Process Model Among Canadian Adults. *Canadian Psychology/Psychologie canadienne*. Advance online publication. January 11. <http://dx.doi.org/10.1037/cap0000270>

Lustria, M. L. A., Noar, S. M., Cortese, J., Van Stee, S. K., Glueckauf, R. L., & Lee, J. 2013. A meta-analysis of web-delivered tailored health behavior change interventions. *Journal of Health Communication*, 18(9), 1039-1069.

McKay, D. L., Berkowitz, J. M., Blumberg, J. B., & Goldberg, J. P. 2004. Communicating cardiovascular disease risk due to elevated homocysteine levels: Using the EPPM to develop print materials. *Health Education & Behavior*, 31(3), 355-371.

National Income Dynamics Study-Coronavirus Rapid Mobile Survey (NIDS-CRAM) 2021, Wave 4 [dataset]. Version Beta3. Cape Town: Allan Gray Orbis Foundation [funding agency]. Cape Town: Southern Africa Labour and Development Research Unit [implementer], 2021. Cape Town: DataFirst [distributor], 2021.

Nino M, Harris C, Drawve G, Fitzpatrick K M. 2021. Race and ethnicity, gender, and age on perceived threats and fear of COVID-19: Evidence from two national data sources, *SSM - Population Health*, Vol13, <https://doi.org/10.1016/j.ssmph.2020.100717>.

Noar, S. M. 2004. A health educator's guide to theories of health behavior. *International Quarterly of Community Health Education*, 24(1), 75-92.

Noar S. M., Grant Harrington, N., Van Stee, S. K., & Shemanski Aldrich, R. 2011. Tailored health communication to change lifestyle behaviors. *American Journal of Lifestyle Medicine*, 5(2), 112-122.

Oyenubi A, Nwosu C., Kollamparambil, U. 2021. The flip-side of risk perception: On the negative effect of risk perception on subjective health during the pandemic. NIDS-CRAM wave 4.

Pijls BG, Jolani S, Atherley A, et al. 2021. Demographic risk factors for COVID-19 infection, severity, ICU admission and death: a meta-analysis of 59 studies, *BMJ Open*;11:e044640. doi: 10.1136/bmjopen-2020-044640

Rimal, R. N. 2001. Perceived risk and self-efficacy as motivators: Understanding individuals' long-term use of health information. *Journal of Communication*, 51, 633-654. doi:10.1111/ j.1460-2466.2001.tb02900.x

Rogers, R. W. 1975. A protection motivation theory of fear appeals and attitude change. *Journal of Psychology*, 91, 93-114.

Rosenstock IM. 1974. Historical origins of the health belief model. *Health Educ Monogr*; 2:328 – 86.

Rubin R. 2021. COVID-19 Vaccines vs Variants—Determining How Much Immunity Is Enough. *JAMA*. 325(13):1241–1243. doi:10.1001/jama.2021.3370

Sallam, M. 2021. "COVID-19 Vaccine Hesitancy Worldwide: A Concise Systematic Review of Vaccine Acceptance Rates" *Vaccines* 9, no. 2: 160. <https://doi.org/10.3390/vaccines9020160>

Shirahmadi S, Seyedzadeh-Sabounchi S, Khazaei S, Bashirian S, Miresmæili AF, Bayat Z, et al. 2020. Fear control and danger control amid COVID-19 dental crisis: Application of the Extended Parallel Process Model. *PLoS ONE* 15(8): e0237490. <https://doi.org/10.1371/journal.pone.0237490>

Slonim, A. B., Roberto, A. J., Downing, C. R., Adams, I. F., Fasano, N. J., Davis-Satterla, L., & Miller, M. A. 2005. Adolescents' knowledge, beliefs, and behaviors regarding hepatitis B: Insights and implications for programs targeting vaccine-preventable diseases. *Journal of Adolescent Health*, 36(3), 178-186.

Smith PJ, Humiston SG, Marcuse EK, Zhao Z, Dorell CG, Howes C, et al. 2011. Parental delay or refusal of vaccine doses, childhood vaccination coverage at 24 months of age, and the Health Belief Model. *Public Health Rep*; 126:Suppl 2 135 – 46.

Taylor A (2021) Blood clot risks: comparing the AstraZeneca vaccine and the contraceptive pill, *The Conversation*. Blood clot risks: comparing the AstraZeneca vaccine and the contraceptive pill ([theconversation.com](https://theconversation.com))

Thompson EL, Vamos CA, Straub DM, Sappenfield WM, Daley EM. 2016. "We've Been Together. We Don't Have It. We're Fine." How Relationship Status Impacts Human Papillomavirus Vaccine Behavior among Young Adult Women. *Womens Health Issues*. 2017 Mar-Apr;27(2):228-236. doi: 10.1016/j.whi.2016.09.011. Epub 2016 Oct 27. PMID: 28277236.

Umberson, D. 1992. Gender, marital status and the social control of health behavior. *Social Science & Medicine*, 34(8), 907–917. [https://doi.org/10.1016/0277-9536\(92\)90259-S](https://doi.org/10.1016/0277-9536(92)90259-S)

- Witte, K. 1992. Putting the fear back into fear appeals: The extended parallel process model. *Communications Monographs*, 59(4), 329-349.
- Witte, K. 1994. Fear control and danger control: A test of the extended parallel process model (EPPM). *Communications Monographs*, 61(2), 113-134.
- Witte, K. 1997. Preventing teen pregnancy through persuasive communications: realities, myths, and the hard-fact truths. *Journal of Community Health*, 22(2), 137-154.
- Witte, K., & Allen, M. 2000. A meta-analysis of fear appeals: Implications for effective public health campaigns. *Health Education & Behavior*, 27(5), 591-615.
- Witte, K., Girma, B., & Girgre, A. 2002. Addressing underlying mechanisms to HIV/AIDS preventive behaviors in Ethiopia. *International Quarterly of Community Health Education*, 21(2), 163-176.
- Wittenberg, M., 2017. Wages and wage inequality in South Africa 1994–2011: part 2–inequality measurement and trends. *South African Journal of Economics*, 85(2), pp.298-318.
- Yang, J., Wu, X., Sasaki, K., & Yamada, Y. 2020. Changing health compliance through message repetition based on the extended parallel process model in the COVID-19 pandemic. *PeerJ*, 8, e10318. <https://doi.org/10.7717/peerj.10318>

## Appendix

**Table A1: Restricted sample logit estimation: Vaccine Hesitancy**

VARIABLES	(1) HBM	(2) EPPM
Risk perception	-0.342** (0.146)	
Efficacy	-1.232*** (0.173)	
Low risk		0.451*** (0.135)
Denial		1.155*** (0.251)
Age	-0.684*** (0.239)	-0.497** (0.204)
Chronic illness	-0.233 (0.179)	-0.319** (0.155)
Covid information	-0.279 (0.225)	-0.385* (0.210)
Log HHincome per capita	-0.0974** (0.0449)	-0.0883** (0.0406)
Education	-0.0375** (0.0191)	-0.0359** (0.0173)
African	-0.846*** (0.223)	-0.689*** (0.205)
NPI behaviour	-0.133** (0.0655)	-0.156*** (0.0564)
Male	-0.138 (0.138)	-0.119 (0.125)
Married	0.0441 (0.141)	-0.0108 (0.125)
Employed	0.251* (0.151)	0.0862 (0.136)
Religious	-0.119 (0.282)	-0.0826 (0.249)
Constant	2.228*** (0.558)	0.819 (0.521)
<b>Wald chi2</b>	<b>128.66***</b>	<b>99.28***</b>
<b>Observations</b>	<b>3,628</b>	<b>3,936</b>

Standard errors in parentheses, \*\*\* p<0.01, \*\* p<0.05, \* p<0.1

**Table A2: Seemingly Unrelated Regression: Health Belief Model (NPI behaviour)**

VARIABLES	(1)	(2)	(3)
	Use sanitiser	facemask	handwash
Risk perception	0.0854*** (0.0170)	0.0645*** (0.0137)	0.0551*** (0.0175)
Efficacy	0.0409* (0.0221)	0.00557 (0.0178)	0.0447** (0.0227)
Age	-0.00476*** (0.000664)	-0.000125 (0.000536)	0.00427*** (0.000685)
Chronic illness	0.0436** (0.0212)	-0.0189 (0.0171)	-0.0129 (0.0219)
Covid information	0.0323 (0.0259)	-0.0137 (0.0209)	0.0378 (0.0267)
Log HHincome per capita	0.00579 (0.00572)	-0.00520 (0.00462)	-0.0102* (0.00591)
Education	0.00424* (0.00252)	0.00844*** (0.00203)	0.000167 (0.00259)
African	0.0235 (0.0218)	0.00249 (0.0176)	0.0688*** (0.0224)
Male	0.0299* (0.0171)	0.0111 (0.0138)	-0.0377** (0.0176)
Married/ Partnered	0.0417** (0.0174)	0.00891 (0.0141)	0.000640 (0.0180)
Urban	0.0412** (0.0199)	0.0801*** (0.0161)	0.0663*** (0.0205)
Employed	0.0552*** (0.0180)	0.0117 (0.0145)	-0.0161 (0.0186)
Religious	0.00718 (0.0292)	0.0368 (0.0236)	0.0708** (0.0301)
Constant	0.502*** (0.0668)	0.613*** (0.0540)	0.205*** (0.0689)
<b>Chi2</b>	<b>237.34***</b>	<b>110.14***</b>	<b>104.27***</b>
<b>Observations</b>	<b>3,504</b>	<b>3,504</b>	<b>3,504</b>
<b>R-squared</b>	<b>0.046</b>	<b>0.026</b>	<b>0.026</b>

Standard errors in parentheses, \*\*\* p<0.01, \*\* p<0.05, \* p<0.1

**Table A3: Seemingly Unrelated Regression: Extended Parallel Process Model (NPI behaviour)**

VARIABLES	(1)	(2)	(3)
	usesanitiser	facemask	handwash
Low risk	-0.0977*** (0.0169)	-0.0679*** (0.0139)	-0.0483*** (0.0176)
Denial (high risk-low efficacy)	-0.102*** (0.0315)	-0.00271 (0.0260)	-0.0815** (0.0328)
Age	-0.00492*** (0.000632)	-0.000391 (0.000520)	0.00361*** (0.000657)
Chronic illness	0.0529*** (0.0204)	-0.00812 (0.0168)	-0.00102 (0.0212)
Covid information	0.0328 (0.0251)	-0.00470 (0.0206)	0.0559** (0.0260)
Log HHincome per capita	0.00998* (0.00550)	-0.00473 (0.00453)	-0.0100* (0.00571)
Education	0.00570** (0.00239)	0.00910*** (0.00197)	-0.000973 (0.00248)
African	0.0176 (0.0209)	0.00818 (0.0172)	0.0533** (0.0217)
Male	0.0289* (0.0164)	0.00182 (0.0135)	-0.0379** (0.0170)
Married/ Partnered	0.0477*** (0.0167)	0.00761 (0.0138)	0.0121 (0.0174)
Urban	0.0366* (0.0191)	0.0731*** (0.0157)	0.0449** (0.0198)
Employed	0.0501*** (0.0172)	0.00642 (0.0142)	-0.00334 (0.0179)
Religious	0.0205 (0.0283)	0.0211 (0.0233)	0.0766*** (0.0294)
Constant	0.595*** (0.0653)	0.697*** (0.0537)	0.338*** (0.0678)
<b>Chi2</b>	<b>197.42***</b>	<b>158.35***</b>	<b>110.47***</b>
<b>Observations</b>	<b>3,791</b>	<b>3,791</b>	<b>3,791</b>
<b>R-squared</b>	<b>0.052</b>	<b>0.025</b>	<b>0.022</b>

**Standard** errors in parentheses, \*\*\* p<0.01, \*\* p<0.05, \* p<0.1

For further information please see [cramsurvey.org](https://cramsurvey.org)